

## POSTER PRESENTATION

## Open Access

# Role of low and high-voltage activated $\text{Ca}^{2+}$ -dependent $\text{K}^{+}$ currents in the control of alpha-motoneuron discharge and its implication in hyperreflexia

Sharmila Venugopal<sup>1</sup>, Thomas M Hamm<sup>1,3</sup>, Ranu Jung<sup>1,2\*</sup>

From Nineteenth Annual Computational Neuroscience Meeting: CNS\*2010  
San Antonio, TX, USA. 24-30 July 2010

Specificity of calcium-activated potassium ( $\text{K}^{+}$ ) currents to different sources of calcium has been noted in many neurons (e.g.<sup>1</sup>). Recently, in spinal alpha-motoneurons ( $\alpha$ -MN), it was shown that the low-voltage activated L-type calcium currents (also known as persistent calcium currents) activate an exclusive subset of small conductance  $\text{K}^{+}$  currents ( $\text{SK}_L$ )<sup>2</sup>. The  $\text{SK}_L$  currents were distinct from the medium after-hyperpolarization (mAHP) producing N/P-type calcium activated  $\text{K}^{+}$  currents ( $\text{SK}_{\text{AHP}}$  currents). The same study further suggested that an enhancement of persistent calcium current often observed after chronic spinalization can in part be due to reduced availability of the  $\text{SK}_L$  channels albeit mAHP remained unchanged. While mAHP has been suggested to be integral in controlling motoneuron firing frequencies and grading L-Ca activation, the role of  $\text{SK}_L$  currents in motoneuron discharge is unknown. The goal of this study is to characterize the influence of  $\text{SK}_{\text{AHP}}$  and  $\text{SK}_L$  currents on motoneuron firing frequencies. Here we test the hypothesis that  $\text{SK}_{\text{AHP}}$  and  $\text{SK}_L$  currents play differential roles in the control of persistent inward currents that are key determinants of motoneuron excitability.


## Methods

The  $\alpha$ -MN is modeled with two compartments (soma and dendrite) using conductance-based Hodgkin-Huxley formalism. The persistent L-Ca and  $\text{SK}_L$  are located in the dendrite along with persistent sodium current. The mAHP causing high-voltage activated  $\text{Ca}^{2+}$  and  $\text{SK}_{\text{AHP}}$

currents are confined to the soma along with action potential causing fast sodium and delayed rectifier  $\text{K}^{+}$  currents. Model simulations are performed using the XPPAUT software.

## Results

The model  $\alpha$ -MN shows counter-clockwise hysteresis in the injected current-frequency (I-f) relationship (Fig. 1, control) as observed in many chronic spinal sacrocaudal motoneurons. This hysteresis is mediated by the dendritic L-Ca and persistent sodium currents (together termed PIC). A selective blockade of somatic  $\text{SK}_{\text{AHP}}$  greatly increases the spike frequencies consistent with experimental findings that mAHP is integral for controlling  $\alpha$ -MN frequencies. On the other hand, eliminating  $\text{SK}_L$  resulted in uncontrolled L-Ca activation with virtually no deactivation of the persistent inward currents



\* Correspondence: Ranu.Jung@asu.edu

<sup>1</sup>Center for Adaptive Neural Systems, Arizona State University, Tempe, AZ-85287, USA

even with large hyperpolarization (self-sustained discharge for  $I_{\text{SOMA}} \leq 0$  does not terminate; compare with  $SK_{\text{AHP}} = 0$  and control traces).

## Conclusions

Chronic spinal cord injury often results in spasticity (hyperreflexia). Intrinsic hyper excitability of  $\alpha$ -MN has been attributed to underlie hyperreflexia. The uncontrolled and abrupt PIC activation due to reduction in  $SK_L$  currents implicates rapid development and sustenance of muscle contraction forces such as during spasms, thus delineating a possible mechanism for  $\alpha$ -MN hyper excitability that could lead to hyperreflexia following injury.

## Acknowledgements

This work was supported by NIH R01-NS054282.

## Author details

<sup>1</sup>Center for Adaptive Neural Systems, Arizona State University, Tempe, AZ-85287, USA. <sup>2</sup>School of Biological and Health Systems Engineering, Arizona State University, Tempe, AZ-85287, USA. <sup>3</sup>Barrow Neurological Institute, Phoenix, AZ-85013, USA.

Published: 20 July 2010

## References

1. Marrion NV, Tavalin SJ: Selective activation of  $Ca^{2+}$ -activated  $K^+$  channels by co-localized  $Ca^{2+}$  channels in hippocampal neurons. *Nature* 1998, **395**:900-905.
2. Li X, Bennett DJ: Apamin-sensitive calcium-activated potassium currents (SK) are activated by persistent calcium currents in rat motoneurons. *J Neurophysiol* 2007, **97**:3314-3330.

doi:10.1186/1471-2202-11-S1-P158

**Cite this article as:** Venugopal *et al.*: Role of low and high-voltage activated  $Ca^{2+}$ -dependent  $K^+$  currents in the control of alpha-motoneuron discharge and its implication in hyperreflexia. *BMC Neuroscience* 2010 **11**(Suppl 1):P158.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

